

(281) 762-6300

COLONOSCOPY PREP

	gister at the location of your procedure as indicated below at least 2-3 vs prior to your procedure. On the day of your procedure,, please arrive at the designated area at
_	·
	Memorial Hermann Ambulatory Surgery Center (ASC) Methodist Sugar Land Hospital Sugar Land Surgical Hospital
arra	cause you will be receiving conscious sedation, please make angements for someone to be responsible for driving you home after procedure.
Jel	ear Liquids permitted the day before your procedure: Broth, Boullion, l-o, Black Coffee, Tea, Apple Juice, Grape Juice, Gatorade, Cranberry ce and Soda.
	MAY NOT have any solid food, Orange Juice, Grapefruit Juice, mato Juice, Pineapple Juice, Milk or Alcohol.
Me	 dications the day of the Procedure If you take Blood Pressure or heart medication, take your medication the morning of your procedure with a small sip of water. All other medication can be taken later in the day after the procedure. For diabetic patients:
	- If you are on blood thinner medication:

Follow Recommended Prep Instructions



SUTAB Colon Prep

SUTABS You will take 2 doses of 12 tablets each. Water must be consumed with each dose of SUTAB and additional water must be consumed after each dose.

PN	
	STEP 1 Open 1 bottle of 12 tablets.
	STEP 2 Fill the provided container with 16 ounces of water (up to the fill line). Swallow each tablet with a sip of water, and drink the entire amount of water over 15 to 20 minutes.
	Tablets not shown actual size.
	IMPORTANT: If you experience preparation-related symptoms (for example, nausea, bloating, or cramping), pause or slow the rate of drinking the additional water until your symptoms diminish.
	Drink additional water
	STEP 3 Approximately 1 hour after the last tablet is ingested, fill the provided container again with 16 ounces of water (up to the fill line), and drink the entire amount over 30 minutes.
	STEP 4 Approximately 30 minutes after finishing the second container of water, fill the provided container with 16 ounces of water (up to the fill line), and drink the entire amount over 30 minutes.
DOSE Two: Ta	ke 12 tablets with water on at

*** NOTHING BY MOUTH AFTER MIDNIGHT ***